



SCIENTIFIC ANNALS of the School of Geology,
Aristotle University of Thessaloniki

SPECIAL VOLUME 102



GREVENA
SIATISTA
GREECE 2014

VIth International
Conference
on Mammoths
and their Relatives

ABSTRACT BOOK

Editors:

Dimitris S. KOSTOPOULOS, Evangelos VLACHOS, and Evangelia TSOUKALA

THESSALONIKI, MAY 2014

Cause and timing of late Early – early Middle Pleistocene European mammoth turnover with special emphasis on Hungary

Attila VIRÁG ✉, and Piroska PAZONYI

During the period between 1.0-0.7 Ma, the climate became drier and colder, consequently the forested areas were rolled back and the steppe vegetation was expanded in Europe. The process begun in the Eastern margin of the continent (e.g. at Taman Peninsula or at the lower Don River area; Tesakov, 2008 and Shchelinsky et al., 2010), and coincided with the first immigration of the steppe mammoth (*Mammuthus trogontherii*), which progressively replaced the indigenous *Mammuthus meridionalis* (often associated with closed habitats). The transition between the two species in Europe was achieved by input from Eastern Asia, either in the form of migrating herds or by gene flow without the long distance movement of individual animals (Lister et al., 2005).

Micromammal assemblages from Russia (Sarkel and Sinyaya Balka) indicate mosaic environment combining closely spaced steppe-like (*Spermophilus*, *Spalax*, *Allocricetus*, *Cricetulus*, *Cricetus*), wooded (*Sorex*, *Apodemus*, *Clethrionomys*), meadow (*Desmana*), and desert (*Plioscirotopoda*) biotopes under conditions of relatively warm climate. At the Taman Peninsula, both *M. meridionalis* and *M. trogontherii* are documented (third molars with low tooth crown and 14-15 plates and third molars with 16-19 plates and higher crown). Van Essen (2011) mentioned some individuals, which show mosaic characteristics (low crown and high plate number) due to the hybridization of the overlapping *M. meridionalis* and *M. trogontherii* populations in such a mosaic environment. At

the same time, *M. trogontherii* was absent from the central and western part of Europe (e.g. Saint-Prest, France, where the large mammalian community indicates an open forest landscape under a temperate wet climate; Guerin et al., 2003).

The older layers of the rich 900 ka micromammal fauna from the Somssich Hill 2 locality (South Hungary) are characterized by the dominance of *Lagurus* and cricetids (mainly *Cricetus runtonensis*), which are indicators of probably the oldest known open steppe vegetation of Hungary. In contrast, mice (*Apodemus*), dormice (*Glis*, *Muscardinus*, *Dryomimus*) and the dominance of voles like *Pitymys*, *Myodes* and *Pliomys* in younger parts of the section are indicative of warmer, more humid climate and more closed (shrubby or forested) habitats (Pazonyi et al., 2013). *M. trogontherii* remains has not been found so far from this period in Hungary.

The 800 ka micromammal fauna of Villány 8 (South Hungary) indicates mosaic environment with the co-occurrence of both steppe (e.g. *Spermophilus*, *Spalax*, *Cricetus*, *Allocricetus*, *Prolagurus*) and forest elements (e.g. *Sorex*, *Apodemus*, *Myodes*) (Jánossy, 1986).

About 800-700 ka, the opening of the vegetation reached Western Europe. The first *M. trogontherii* remains in the central and western part of the continent can be dated to the beginning of this period, nevertheless, contemporaneously *M. meridionalis* populations were also




Fig. 1. *Mammuthus trogontherii* upper left third molar (Hungarian Natural History Museum, V.72.116) from Üröm Hill (North-Hungary) from lingual (A) and occlusal (B) view. C, posterior view of a plate in midcrown area. The arrows on A indicate well-developed cement furrows.

documented in the region (e.g. from Győrújfalú in Hungary or from the Ponte Galeria Formation in Italy, and from Dorn-Dürkheim 3 in Germany; Palombo & Ferretti, 2005 and van Essen, 2011). Despite the slight recovery of closed vegetation, *M. meridionalis* almost entirely disappeared from Europe by the end of this period. Only the remains of hybrid individuals support, that the last representatives of the species had to be present in some refugia at that time. Such mosaic specimen (an upper third molar with a reconstructed plate number of 17 or even 18 and relatively high crown but with plesiomorph enamel, thicker than 3.0 mm, see Fig. 1. and Virág, 2009 for details) was found at the 800-700 ka locality, called Üröm Hill (North Hungary). Similar specimens turned out from Campo del Conte in Italy (although Palombo et al., 2003 identified the molar as *M. meridionalis*) or from Voigtstedt and Edersleben in Germany (van Essen, 2011).

The fauna of Üröm Hill mainly contain hygrophilous taxa (like *Desmana thermalis*, *Mimomys savini* or *Hippopotamus antiquus*) and the micromammal fauna of the contemporaneous Kövesvára locality (North Hungary) unequivocally indicate warm, humid climate and closed, forested vegetation with the dominance of dormice (*Glis*, *Muscardinus*) and forest voles (*Myodes*) (Jánossy, 1986). Similar warm-temperate conditions with mainly forest elements and scanty open landscape indicators (such as *Spermophilus*) are typical for Voigtstedt (Maul & Parfitt, 2010).

By 600 ka, only *M. trogontherii* occurred in Europe, as at Süssenborn (Germany), the type locality of the species (Lister et al., 2005). The extinction of the *M. meridionalis* populations was most plausibly triggered not solely by the immigration of the steppe-adapted, grazer *M. trogontherii* from Eastern Asia, but the arrival of the browser or mixed-feeder *Elephas antiquus* from Africa through the Levant, which happened broadly 900-800 ka (Albayrak & Lister, 2012). The earliest occurrence of the latter species in Hungary is most plausibly Győrújfalú. Following this period, *E. antiquus* has unfortunately very scanty record in Hungary.

This project was supported by the OTKA K 104506 grant.

References

- Albayrak, E., Lister, A.M., 2012. Dental remains of fossil elephants from Turkey. *Quaternary International* 276-277, 198-211.
- Guérin, C., Dewolf, Y., Lautridou, J., 2003. Revision of a famous site: Saint-Prest. *Geobios* 36(1), 55-82.
- Jánossy, D., 1986. Pleistocene Vertebrate Faunas of Hungary. *Developments in Palaeontology and Stratigraphy* 8, 1-208.
- Lister, A.M., Sher, A.V., van Essen, H., Wei, G., 2005. The pattern and process of mammoth evolution in Eurasia. *Quaternary International* 126-128, 49-64.
- Maul, L.C., Parfitt, S.A., 2010. Micromammals from the 1995 Mammoth Excavation at West Runton, Norfolk, UK: Morphometric data, biostratigraphy and taxonomic reappraisal. *Quaternary International* 228(1-2), 91-115.
- Palombo, M.R., Ferretti, M.P., 2005. Elephant fossil record from Italy: knowledge, problems, and perspectives. *Quaternary International* 126-128, 107-136.
- Palombo, M.R., Magri, D., Molinaro, A., Pisano, V., 2003. The Pleistocene sequence of Campo del Conte (Lower Sacco Valley, Southern Lazio). *Geologica Romana* 36, 289-309.
- Pazonyi, P., Mészáros, L., Szentesi, Z., Gasparik, M., Virág, A., 2013. Preliminary results of the palaeontological investigations of the late Early Pleistocene Somssich Hill 2 locality (South Hungary). 14th Congress of Regional Committee on Mediterranean Neogene Stratigraphy, Book of Abstracts of the RCMNS 2013, Istanbul Technical University, Istanbul, 270.
- Shchelinsky, V.E., Dodonov, V.S., Baigusheva, V.S., Kulakov, S.A., Simakova, A.N., Tesakov, A.S., Titov, V.V., 2010. Early Palaeolithic sites on the Taman Peninsula (Southern Azov Sea region, Russia): Bogatyri/Sinyaya Balka and Rodniki. *Quaternary International* 223-224, 28-35.
- Tesakov, A.S., 2008. Early Pleistocene mammalian fauna of Sarkel (Lower Don River area, Russia): mole voles (Ellobiusini, Arvicolinae, Rodentia). *Russian Journal of Theriology* 7(2), 81-88.
- Van Essen, H., 2011. Tracing transitions. An overview of the evolution and migrations of the genus *Mammuthus* Brookes, 1828 (Mammalia, Proboscidea). PhD Theses. University of Leiden, Faculty of Archaeology, 1-270.
- Virág, A., 2009. Revision of the "*Archidiskodon meridionalis ürömensis*" Vörös, 1979 in the context of the mammoth evolution in Eurasia. *Central European Geology* 52(3-4), 405-419.

✉ myodes.glareolus@gmail.com



Citation:

Virág, A., Pazonyi, P., 2014. Cause and timing of late Early – early Middle Pleistocene European mammoth turnover with special emphasis on Hungary. *Abstract Book of the VIth International Conference on Mammoths and their Relatives. S.A.S.G., Special Volume 102: 211-212*